Journal of Mathematical Extension
Vol. 19, No. 6 (2025) (5) 1-13
ISSN: 1735-8299
URL: http://doi.org/10.30495/JME.2025.3353
Original Research Paper

Generalized Volterra-Type Operators from Hardy Space into Iterated Weighted-Type Spaces

E. Abbasi
Islamic Azad University, Mahabad, Iran

D. Molaei *
Islamic Azad University, Urmia, Iran

A. Ebrahimi
Islamic Azad University, Mahabad, Iran

Abstract. Let $H(\mathbb{D})$ be the set of analytic functions on $\mathbb{D}$ and for $1 \leq p \leq \infty$, H^p be the Hardy space. For $m \in \mathbb{N}$ suppose that I^m be mth iteration. Let $\overrightarrow{g} = (g_0, \cdots, g_{m-1})$ where $\{g_i\}_{i=0}^{m-1} \subset H(\mathbb{D})$ and $I(f) = \int_0^z f(w)dw$. If I^m for $m \in \mathbb{N}$ be the mth iteration, then the generalized Volterra-type operators $I_{\overrightarrow{g}}^m$ on $H(\mathbb{D})$ is defined as follows

$$I_{\overrightarrow{g}}^m(f) = I^m(\sum_{i=0}^{m-1} f^{(i)} g_i).$$

In this paper, we investigate boundedness and compactness of generalized Volterra-type operators from Hardy space into iterated weighted-type spaces, $V_n = \{f \in H(\mathbb{D}) : \ \sup_{z\in\mathbb{D}}(1-|z|^2)|f^{(n)}(z)| < \infty\}$.

AMS Subject Classification: 47G10; 30H10; 30H05; 30H99
Keywords and Phrases: Boundedness, Compactness, Generalized Volterra-type operators, Hardy space, Iterated weighted-type spaces

Received: May 2025; Accepted: November 2025
*Corresponding Author

1 Introduction

Let $\mathbb{D}$ be an open unit disc in the complex $\mathbb{C}$ and $H(\mathbb{D})$ be the set of analytic functions on $\mathbb{D}$. For $1 \leq p < \infty$, the Hardy space H^p consists of all analytic functions $f \in H(\mathbb{D})$ such that

$$\|f\|_{H^p} = \sup_{0<r<1} \Big(\frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\theta})|^p d\theta\Big)^{\frac{1}{p}} < \infty.$$

Also H^∞ is the space of bounded analytic functions on $\mathbb{D}$ with the norm $\|f\|_{H^\infty} = \sup_{z\in\mathbb{D}} |f(z)|$. More information about such spaces can be found in [9].

Another space used in this paper is nth weighted-type space. Let μ be a weight (continuous and positive function on $\mathbb{D}$) and $n \in \mathbb{N}_0$. The nth weighted-type space V_n^μ, consists of all analytic functions $f \in H(\mathbb{D})$ such that $b_{V_n^\mu}(f) = \sup_{z\in\mathbb{D}} \mu(z)|f^{(n)}(z)| < \infty$. This space is a Banach with the following norm

$$\|f\|_{V_n^\mu} = \sum_{i=0}^{n-1} \mid f^{(i)}(0) \mid + b_{V_n^\mu}(f) < \infty.$$

For $\alpha > 0$ and $\mu(z) = (1 - |z|^2)^\alpha$, we use V_n^α, V_n and $\|\cdot\|_n$ instead of V_n^μ, V_n^1 and $\|\cdot\|_{V_n^1}$. The space V_n^α contains a large class of analytic functions. For example when $\alpha > 0$, $V_0^\alpha = \mathcal{A}^{-\alpha}$(growth space), $V_1^\alpha = \mathcal{B}^\alpha$ (Bloch type space), $V_2^\alpha = \mathcal{Z}^\alpha$ (Zygmund type space), $V_1 = \mathcal{B}$ (classic Bloch space) and $V_2 = \mathcal{Z}$ (classic Zygmund space). The space V_n is called iterated weighted-type space. In [7] Colonna *et al.* considered iterated weighted-type spaces and obtained some properties for these spaces, especially they showed

$$\cdots \subset V_{n+1} \subset V_n \subset \cdots \subset V_3 \subset \mathcal{Z} \subset H^\infty \subset \mathcal{B} \subset \mathcal{A}^{-1}.$$

The closed subspace of V_n^μ containing of all $f \in V_n^\mu$ such that $\lim_{|z|\to 1} \mu(z)|f^{(n)}(z)| = 0$ is denoted by $V_{n,0}^\mu$ and is called little nth weighted-type space. For more information about (little) nth weighted-type spaces, see [1, 2, 4, 7, 12, 14].

Let $m \in \mathbb{N}$, $I^m(f) = \int_0^z \int_0^{z_1} \cdots \int_0^{z_{m-1}} f(z)dzdz_1 \cdots dz_{m-1}$, and $\overrightarrow{g} = (g_0, g_1, \cdots, g_{m-1})$ where $\{g_i\}_{i=0}^{m-1} \subset H(\mathbb{D})$. The generalized Volterra-type operator on $H(\mathbb{D})$ defined as follows

$$I_{\overrightarrow{g}}^m(f) = I^m(\sum_{i=0}^{m-1} f^{(i)} g_i).$$

For $m = 1$ and $g_0 = g'$, we get Volterra type operator $(J_g f)(z) = \int_0^z f(w)g'(w)dw$ and when $m = 1$ and $g_0 = 1$, we obtain the classic Volterra operator $(If)(z) = \int_0^z f(w)dw$. Also if we set $g_i = a_{m-i-1}g^{(m-i)}$ $(0 \leq i \leq m-1)$, where $g \in H(\mathbb{D})$ and $\overrightarrow{a} = (a_0, \cdots, a_{m-1}) \in \mathbb{C}^m$, we have generalized integration operator $I_{g,\overrightarrow{a}}^m$ defined by Chalmoukis in [6].

Since

$$I_{\overrightarrow{g}}^m(f) = I^m(\sum_{i=0}^{m-1} f^{(i)} g_i) = \sum_{i=0}^{m-1} I^m(f^{(i)} g_i) := \sum_{i=0}^{m-1} I_{g_i}^{m,i}(f),$$

so for considering properties of operator $I_{\overrightarrow{g}}^m$, firstly we investigate properties of operators $I_{g_i}^{m,i}$ where $0 \leq i \leq m-1$.

Chalmoukis in [6] considered boundedness and compactness of $I_{g,\overrightarrow{a}}^m : H^p \to H^q$, where $(0 < q < p < \infty)$ and posed a conjecture that g must be in $H^{\frac{pq}{q-p}}$. Yang *et al.* provided a positive answer to the aforementioned conjecture in [13]. Arroussi *et al.* investigated boundedness and compactness of $I_{\overrightarrow{g}}^m : A^p \to A^q$, where A^p is Bergman space. They extended Chalmoukis result to Bergman spaces and showed that the Bergman space version of Chalmoukis conjecture is true (see [5]). Also some authors characterized boundedness and compactness of generalized integration operators among some other analytic function spaces [8, 10]. In [14], Zhu investigated Bloch-type spaces and uncovered numerous properties associated with these spaces. Later, Stevi expanded on this concept by generalizing Bloch-type spaces and introducing the nth weighted-type spaces, as detailed in [11, 12]. In recent years, extensive research has been conducted on such spaces, with one of the most notable references in this area being [7].

In this paper, firstly we investigate boundedness and compactness of the operators $I_{g_i}^{m,i} (0 \leq i \leq m)$ from Hardy space into iterated weighted-type spaces and we find some characterizations for boundedness and

compactness of such operators. Then we consider boundedness and compactness of the operator $I_{\overrightarrow{g}}^{m} : H^p \to V_n$ and we show that the operator $I_{\overrightarrow{g}}^{m} : H^p \to V_n$ is bounded (compact) if and only if each operator $I_{g_i}^{m,i} : H^p \to V_n (0 \leq i \leq m-1)$ is bounded (compact).

In this work, we shall use the notation $A \preceq B$ to mean that for some $c > 0$, $A \leq c\, B$, whereas $A \asymp B$ means $A \preceq B$ and $B \preceq A$.

2 Boundedness and Compactness of Operator $I_{g_i}^{m,i} : H^p \to V_n$

In this section, we investigate boundedness and compactness of the operators $I_{g_i}^{m,i}$ from Hardy space $H^p (1 \leq p \leq \infty)$ into iterated weighted-type spaces and we obtain some characterizations for boundedness and compactness of such operators. We begin with the following lemma.

Lemma 2.1. *Let $n, k \in \mathbb{N}$. For any $f \in V_n$, we have*

$$\|f\|_n \asymp \sum_{i=0}^{n+k-2} |f^{(i)}(0)| + \sup_{z\in\mathbb{D}} (1-|z|^2)^k |f^{(n+k-1)}(z)|.$$

Proof. For $n = 1$, $V_1 = \mathcal{B}$. So by using Proposition 8 in[14], we get

$$\|f\|_1 = \|f\|_{\mathcal{B}} \asymp \sum_{i=0}^{k-1} |f^{(i)}(0)| + \sup_{z\in\mathbb{D}} (1-|z|^2)^k |f^{(k)}(z)|.$$

For any $n \in \mathbb{N}$, $V_n \subset V_{n-1}$ [6, Proposition 2.1]. Hence, for any $f \in V_n$, $f^{(n-1)} \in \mathcal{B}$. By replacing f with $f^{(n-1)}$ in the above equation, we obtain

$$|f^{(n-1)}(0)| + \sup(1-|z|^2)|f^{(n)}(z) \mid \asymp \sum_{i=n-1}^{n+k-2} |f^{(i)}(0)| + \sup_{z\in\mathbb{D}} (1-|z|^2)^k |f^{(n+k-1)}(z)|.$$

Hence,

$$\begin{aligned}\|f\|_n &= \sum_{i=0}^{n-2}|f^{(i)}(0)| + |f^{(n-1)}(0)| + \sup_{z\in\mathbb{D}}(1-|\ z\ |^2)\ |\ f^{(n)}(z)\ | \\ &\asymp \sum_{i=0}^{n-2}|f^{(i)}(0)| + \sum_{i=n-1}^{n+k-2}|f^{(i)}(0)| + \sup_{z\in\mathbb{D}}(1-|\ z\ |^2)^k\ |\ f^{(n+k-1)}(z)\ | \\ &= \sum_{i=0}^{n+k-2}|f^{(i)}(0)| + \sup_{z\in\mathbb{D}}(1-|\ z\ |^2)^k|f^{(n+k-1)}(z)|.\end{aligned}$$

The proof is complete. □

Let $m \geq n$. It is clear $\left(I_{\overrightarrow{g}}^m(f)\right)^{(i)}(0) = \left(I_g^{m,i}(f)\right)^{(i)}(0) = 0$, when $0 \leq i \leq m-1$, so for any $f \in H(\mathbb{D})$, by using Lemma 2.1, we have

$$\begin{aligned}\|I_g^{m,i}(f)\|_n &\asymp \sum_{k=0}^{m-1}\left|\left(I_g^{m,i}(f)\right)^{(k)}(0)\right| + \sup_{z\in\mathbb{D}}(1-|z|^2)^{m-n+1}\left|f^{(i)}g(z)\right| \quad (1)\\ &= \sup_{z\in\mathbb{D}}(1-|z|^2)^{m-n+1}\left|f^{(i)}g(z)\right|.\end{aligned}$$

Lemma 2.2. *Let $1 \leq p \leq \infty$. Then for any $f \in H^p$ and $k \in \mathbb{N}_0$,*

$$\sup_{z\in\mathbb{D}}(1-|z|^2)^{k+\frac{1}{p}}|f^{(k)}(z)| \preceq \|f\|_{H^p}.$$

Proof. From Proposition 5.1.2 of [14], we have $H^\infty \subset \mathcal{B}$ and for any $f \in H^\infty$

$$\sup_{z\in\mathbb{D}}(1-|z|^2)|f'(z)| \leq \|f\|_{H^\infty}.$$

Applying Lemma 2.1, for each $k \in \mathbb{N}_0$ and $f \in H^\infty$, we obtain

$$\sup_{z\in\mathbb{D}}(1-|z|^2)^k|f^{(k)}(z)| \leq \|f\|_1 \leq 2\|f\|_{H^\infty} \preceq \|f\|_{H^\infty}.$$

Similar results for $1 \leq p < \infty$ follow easily using results from [9]. □

Theorem 2.3. *Let $1 \leq p \leq \infty$, $m, n \in \mathbb{N}$ such that $m \geq n$ and $g_i \in H(\mathbb{D})$. Then for each $0 \leq i \leq m$, the operator $I_{g_i}^{m,i} : H^p \to V_n$ is bounded*

if and only if $\sup_{z\in\mathbb{D}}(1-|z|^2)^{m-n-i-\frac{1}{p}+1}|g_i(z)|<\infty$. *Moreover, in this case*

$$\|I_{g_i}^{m,i}\| \asymp \sup_{z\in\mathbb{D}}(1-|z|^2)^{m-n-i-\frac{1}{p}+1}|g_i(z)|.$$

Proof. Let the operator $I_{g_i}^{m,i}: H^p \to V_n$ be bounded and $f_{i,w}(z)=\frac{(1-|w|^2)^i}{(1-\overline{w}z)^{i+\frac{1}{p}}}$, where $0\leq i\leq m$ and $w\in\mathbb{D}-\{0\}$. For $p=\infty$,

$$\sup_{z\in\mathbb{D}}|f_{i,w}(z)| = \sup_{z\in\mathbb{D}}\left|\frac{1-|w|^2}{1-\overline{w}z}\right|^i \leq \sup_{z\in\mathbb{D}}\frac{(1-|w|^2)^i}{(1-|\overline{w}|)^i}=2^i,$$

so, $\sup_{w\in\mathbb{D}}\|f_{i,w}\|_{H^\infty}\leq 2^i$ and when $1\leq p<\infty$, from Lemma 2 [11], there exists positive constant $C_{i,p}$ such that $\sup_{w\in\mathbb{D}}\|f_{i,w}\|_{H^p}<C_{i,p}$. Applying (1) for $f_{i,w}$, we have

$$\begin{aligned}\|I_{g_i}^{m,i}(f_{i,w})\|_n &\asymp \sup_{z\in\mathbb{D}}(1-|z|^2)^{m-n+1}|f_{i,w}^{(i)}(z)g_i(z)|\\ &\geq (1-|w|^2)^{m-n+1}|f_{i,w}^{(i)}(w)||g_i(w)|\\ &= |\overline{w}|^i\prod_{l=0}^{i-1}(i+\frac{1}{p}+l)(1-|w|^2)^{m-n-i-\frac{1}{p}+1}|g_i(w)|.\end{aligned}$$

Therefore,

$$\begin{aligned}\sup_{w\in\mathbb{D}}(1-|w|^2)^{m-n-i-\frac{1}{p}+1}|g_i(w)| \preceq \|I_{g_i}^{m,i}(f_{i,w})\|_n &\leq \|I_{g_i}^{m,i}\|\sup_{w\in\mathbb{D}}\|f_{i,w}\|_{H^p}\\ &\leq C_{i,p}\|I_{g_i}^{m,i}\|.\end{aligned}$$

Conversely, we assume that $\sup_{z\in\mathbb{D}}(1-|z|^2)^{m-n-i-\frac{1}{p}+1}|g_i(z)|<\infty$, by using (1), Lemmas 2.1 and 2.2, for any $f\in H^p$, we obtain

$$\begin{aligned}\|I_{g_i}^{m,i}(f)\|_n &\asymp \sup_{z\in\mathbb{D}}(1-|z|^2)^{m-n+1}|f^{(i)}(z)g_i(z)|\\ &\leq \sup_{z\in\mathbb{D}}(1-|z|^2)^{i+\frac{1}{p}}|f^{(i)}(z)|\sup_{z\in\mathbb{D}}(1-|z|^2)^{m-n-i-\frac{1}{p}+1}|g_i(z)|\\ &\preceq \|f\|_{H^p}\sup_{z\in\mathbb{D}}(1-|z|^2)^{m-n-i-\frac{1}{p}+1}|g_i(z)|.\end{aligned}$$

Therefore, the operator $I_{g_i}^{m,i} : H^p \to V_n$ is bounded and $\|I_{g_i}^{m,i}\| \preceq \sup_{z\in\mathbb{D}}(1-|z|^2)^{m-n-i-\frac{1}{p}+1}|g_i(z)|$. The proof is complete. $\square$

To investigate compactness of operators $I_g^{m,i} : H^p \to V_n (0 \leq i \leq m)$, we need the following lemma, since the proof of it is similar to the proof of [7, proposition 3.11], so it is omitted.

Lemma 2.4. *Let $1 \leq p \leq \infty$, $m, n \in \mathbb{N}$, $\{g_i\}_{i=0}^{m-1} \subset H(\mathbb{D})$ and $T = I_{\vec{g}}^m$ or $I_{g_i}^{m,i} (0 \leq i \leq m)$. Then the bounded operator $T : H^p \to V_n$ is compact if and only if for any bounded sequence $\{f_k\}$ in H^p which converges to zero uniformly on compact subsets of $\mathbb{D}$, $\lim_{k\to\infty} \|T(f_k)\|_n = 0$.*

Theorem 2.5. *Let $1 \leq p \leq \infty$, $m, n \in \mathbb{N}$ such that $m \geq n$ and $g_i \in H(\mathbb{D}) (0 \leq i \leq m)$. Then for each $0 \leq i \leq m$, the operator $I_{g_i}^{m,i} : H^p \to V_n$ is compact if and only if $\lim_{|z|\to 1}(1-|z|^2)^{m-n-i-\frac{1}{p}+1}|g_i(z)| = 0$.*

Proof. Let the operator $I_{g_i}^{m,i} : H^p \to V_n$ be compact. For any $0 \leq i \leq m$ and $w \in \mathbb{D} - \{0\}$ the functions $f_{i,w}(z) = \frac{(1-|w|^2)^i}{(1-\overline{w}z)^{i+\frac{1}{p}}}$ are bounded and converge uniformly to zero on compact subsets of $\mathbb{D}$ when $|w|$ tends to 1, so by applying Lemma 2.4 $\lim_{|w|\to 1} \|I_{g_i}^{m,i}(f_{i,w})\|_n = 0$. Now by using (1), we obtain

$$\begin{aligned}\|I_{g_i}^{m,i}(f_{i,w})\|_n &\asymp (1-|z|^2)^{m-n+1}|f_{i,w}^{(i)}(z)||g_i(z)| \\ &\succeq (1-|w|^2)^{m-n-i-\frac{1}{p}+1}|\overline{w}|^i|g_i(w)|.\end{aligned}$$

In the above inequality, let $|w| \to 1$. Then, we obtain $\lim_{|w|\to 1}(1-|w|^2)^{m-n-i-\frac{1}{p}+1}|g_i(w)| = 0$.

Conversely, suppose that $\lim_{|z|\to 1}(1-|z|^2)^{m-n-i-\frac{1}{p}+1}|g_i(z)| = 0$. Hence, for any $\varepsilon > 0$ there exists $0 < \delta < 1$ such that for each $\delta < |z| < 1$,

$$(1-|z|^2)^{m-n-i-\frac{1}{p}+1}|g_i(z)| < \varepsilon. \tag{2}$$

Now by using (1) and Lemma 2.1, for any bounded sequence $\{f_k\} \subset H^p$

which converges uniformly to zero on compact subsets of $\mathbb{D}$, we obtain

$$\begin{aligned}\|I_{g_i}^{m,i}(f_k)\|_n &\asymp \sup_{z\in\mathbb{D}}(1-|z|^2)^{m-n+1}|f_k^{(i)}(z)||g_i(z)|\\ &\le \sup_{|z|\le\delta}(1-|z|^2)^{m-n+1}|f_k^{(i)}(z)||g_i(z)|\\ &\quad+\sup_{\delta<|z|<1}(1-|z|^2)^{m-n+1}|f_k^{(i)}(z)||g_i(z)|\\ &\le \sup_{|z|\le\delta}(1-|z|^2)^{i+\frac{1}{p}}|f_k^{(i)}(z)|\sup_{|z|\le\delta}(1-|z|^2)^{m-n-i-\frac{1}{p}+1}|g_i(z)|\\ &\quad+\sup_{\delta<|z|<1}(1-|z|^2)^{i+\frac{1}{p}}|f_k^{(i)}|\sup_{\delta<|z|<1}(1-|z|^2)^{m-n-i-\frac{1}{p}+1}|g_i(z)|\\ &=: X_1+X_2.\end{aligned}$$

By using Cauchy's estimates, for any $i\in\mathbb{N}_0$, the sequence $\{f_k^{(i)}\}$ converges uniformly to zero on compact subsets of $\mathbb{D}$, therefore

$$\lim_{k\to\infty}X_1\le\sup_{|z|\le\delta}(1-|z|^2)^{m-n-i-\frac{1}{p}+1}|g_i(z)|\lim_{k\to\infty}\sup_{|z|\le\delta}|f_k^{(i)}(z)|=0.$$

Also applying Lemma 2.2 and (2), we get

$$\begin{aligned}\lim_{k\to\infty}X_2 &\preceq \lim_{k\to\infty}\|f_k\|_{H^p}\sup_{\delta<|z|<1}(1-|z|^2)^{m-n-i-\frac{1}{p}+1}|g_i(z)|\\ &\le \varepsilon\sup_{k\in\mathbb{N}}\|f_k\|_{H^p}.\end{aligned}$$

So, $\lim_{k\to\infty}\|I_{g_i}^{m,i}(f_k)\|_n=0$. By using Lemmas 2.4, the operator $I_{g_i}^{m,i}:H^p\to V_n$ is compact. The proof is complete. □

Putting $n=1$ and $n=2$ in Theorems 2.3 and 2.5, we get the following corollaries.

Corollary 2.6. *Let $1\le p\le\infty$, $m\in\mathbb{N}$ and $g_i\in H(\mathbb{D})$. Then for each $0\le i\le m$, the operator $I_{g_i}^{m,i}:H^p\to\mathcal{B}$ is bounded (compact) if and only if $g_i\in V_0^{m-i-\frac{1}{p}}$ ($g_i\in V_{0,0}^{m-i-\frac{1}{p}}$).*

Corollary 2.7. *Let $1\le p\le\infty$, $m\in\mathbb{N}$ such that $m\ge 2$ and $g_i\in H(\mathbb{D})$. Then for each $0\le i\le m$, the operator $I_{g_i}^{m,i}:H^p\to\mathcal{Z}$ is bounded (compact) if and only if $g_i\in V_0^{m-i-\frac{1}{p}-1}$ ($g_i\in V_{0,0}^{m-i-\frac{1}{p}-1}$).*

3 Boundedness and Compactness of Operator $I^m_{\overrightarrow{g}} : H^p \to V_n$

In this section, we will consider boundedness and compactness of the operator $I^m_{\overrightarrow{g}}$ from Hardy spaces into iterated weighted-type spaces. Especially we show that if the operator $I^m_{\overrightarrow{g}} : H^p \to V_n$ is bounded (compact) then for each $0 \leq i \leq m-1$, the operator $I^{m,i}_{g_i} : H^p \to V_n$ is bounded (compact). For this purpose, we need the following lemma which comes from [2, Lemma 2.5] and [3, Lemma 2.3].

Lemma 3.1. *Let $1 \leq p \leq \infty$. For any $0 \neq \xi \in \mathbb{D}$ and $i \in \{0, 1, \cdots, n\}$, there exists a function $v_{i,\xi} \in H^p$ with the following conditions:*

a) $v_{i,\xi}(z) = \sum_{j=1}^{n+1} c^i_j f_{i,\xi}(z)$, where $f_{i,\xi}(z) = \frac{(1-|\xi|^2)^i}{(1-\bar{\xi}z)^{i+\frac{1}{p}}}$ and c^i_j is independent of choice ξ.

b) $\sup_{\xi\in\mathbb{D}} \|v_{i,\xi}\|_{H^p} < \infty$ and

$$v^{(k)}_{i,\xi}(\xi) = \begin{cases} \frac{\bar{\xi}^i}{(1-|\xi|^2)^{i+\frac{1}{p}}}, & k = i, \\ 0, & k \neq i. \end{cases}$$

c) For any sequence $\{\xi_k\} \subset \mathbb{D}$ such that $\lim_{k\to\infty} |\xi_k| = 1$, the sequence $\{v_{i,\xi_k}\}$ converges to zero uniformly on compact subsets of $\mathbb{D}$.

Let $m \geq n$. For any $f \in H(\mathbb{D})$, applying Lemma 2.1, we get

$$\|I^m_{\overrightarrow{g}}(f)\|_n \asymp \sum_{k=0}^{m-1} \left|\left(I^m_{\overrightarrow{g}}(f)\right)^{(k)}(0)\right| + \sup_{z\in\mathbb{D}} (1-|z|^2)^{m-n+1} \left|\left(I^m_{\overrightarrow{g}}(f)\right)^{(m)}\right| \tag{3}$$

$$= \sup_{z\in\mathbb{D}} (1-|z|^2)^{m-n+1} \left|\sum_{j=0}^{m-1} f^{(j)}(z) g_j(z)\right|.$$

Theorem 3.2. *Let $1 \leq p \leq \infty$ and $m, n \in \mathbb{N}$ such that $m \geq n$. Then the following conditions are equivalent:*

a) The operator $I^m_{\overrightarrow{g}} : H^p \to V_n$ *is bounded.*

b) For each $0 \leq i \leq m-1$, *the operator* $I^{m,i}_{g_i} : H^p \to V_n$ *is bounded.*

c) For each $0 \leq i \leq m-1$, $\sup_{z\in\mathbb{D}}(1-|z|^2)^{m-n-i-\frac{1}{p}+1}|g_i(z)| < \infty$.

Proof. (b)⇒ (a) Since $I^m_{\overrightarrow{g}} = \sum_{i=0}^{m-1} I^{m,i}_{g_i}$ and all operators $I^{m,i}_{g_i} : H^p \to V_n (0 \leq i \leq m-1)$ are bounded, hence the operator $I^m_{\overrightarrow{g}} : H^p \to V_n$ is bounded.

(b)⇔ (c) Theorem 2.3.

(a)⇒ (c) Let the operator $I^m_{\overrightarrow{g}} : H^p \to V_n$ be bounded. For each $i \in \{0, \cdots, m-1\}$ and $\xi \in \mathbb{D} - \{0\}$, let $v_{i,\xi}$ be function found in Lemma 3.1, by using (3), we have

$$\begin{aligned} \|I^m_{\overrightarrow{g}}(v_{i,\xi})\|_{V_n} &\asymp \sup_{z\in\mathbb{D}}(1-|z|^2)^{m-n+1}|\sum_{j=0}^{m-1} v^{(j)}_{i,\xi}(z)g_j(z)| \qquad (4)\\ &\geq (1-|\xi|^2)^{m-n+1}|\sum_{j=0}^{m-1} v^{(j)}_{i,\xi}(\xi)g_j(\xi)|\\ &\geq (1-|\xi|^2)^{m-n+1}|v^{(s)}_{i,\xi}(\xi)||g_i(\xi)|\\ &= (1-|\xi|^2)^{m-n+1}\frac{|\bar{\xi}|^i}{(1-|\xi|^2)^{i+\frac{1}{p}}}|g_i(\xi)|\\ &= |\bar{\xi}|^i(1-|\xi|^2)^{m-n-i-\frac{1}{p}+1}|g_i(\xi)|. \end{aligned}$$

Applying Lemma 3.1(b), we get

$$\sup_{\xi\in\mathbb{D}}(1-|\xi|^2)^{m-n-i-\frac{1}{p}+1}|g_i(\xi)| \leq \|I^m_{\overrightarrow{g}}(v_{i,\xi})\|_{V_n} \leq \|I^m_{\overrightarrow{g}}\| \sup_{\xi\in\mathbb{D}} \|v_{i,\xi}\|_{H^p} < \infty.$$

The proof is complete. □

Theorem 3.3. *Let* $1 \leq p \leq \infty$ *and* $m, n \in \mathbb{N}$ *such that* $m \geq n$. *Then the following conditions are equivalent:*

a) The operator $I^m_{\overrightarrow{g}} : H^p \to V_n$ *is compact.*

b) For each $0 \leq i \leq m-1$, *the operator* $I^{m,i}_{g_i} : H^p \to V_n$ *is compact.*

c) For each $0 \leq i \leq m-1$, $\lim_{|z|\to 1}(1-|z|^2)^{m-n-i-\frac{1}{p}+1}|g_i(z)| = 0$.

Proof. (b)⇒ (a) It is clear when for each $0 \leq i \leq m-1$, the operator $I^{m,i}_{g_i} : H^p \to V_n$ is compact, then $I^m_{\overrightarrow{g}} = \sum_{i=0}^{m-1} I^{m,i}_{g_i}$ is compact.

(b)⇔ (c) Theorem 2.5.

(a)⇒ (c) Assume that the operator $I^m_{\overrightarrow{g}} : H^p \to V_n$ is compact. For each $i \in \{0, \cdots, m-1\}$ and $\xi \in \mathbb{D} - \{0\}$, the sequence $\{v_{i,\xi}\}$ is bounded and converges to zero uniformly on compact subsets of $\mathbb{D}$ when $|\xi|$ tends to 1 (Lemma 3.1), so by using Lemma 2.4, $\lim_{|\xi|\to 1} \|I^m_{\overrightarrow{g}}(v_{i,\xi})\|_n = 0$. Now it is enough to let $|\xi|$ tends to 1 in the inequality (4), therefore $\lim_{|\xi|\to 1}(1-|\xi|^2)^{m-n-i-\frac{1}{p}+1}|g_i(\xi)| = 0$. The proof is complete. □

Let $m \in \mathbb{N}$, $\overline{a} = (a_0, \cdots, a_{m-1}) \in \mathbb{C}^m$ and $g \in H(\mathbb{D})$. Applying Theorems 3.2 and 3.3, we obtain similar results for the Chalmoukis operator

$$I^m_{g,\overline{a}} f(z) = I^m \left(f g^{(n)} + a_1 f' g^{(n-1)} + \cdots + a_{n-1} f^{(n-1)} g' \right)$$

acting from the Hardy space into iterated weighted-type spaces.

Corollary 3.4. *Let* $1 \leq p \leq \infty$, $m \in \mathbb{N}$, $\overline{a} = (a_0, \cdots, a_{m-1}) \in \mathbb{C}^m$ *and* $g \in H(\mathbb{D})$. *Then*

a) the operator $I^m_{g,\overline{a}} : H^p \to V_n$ *is bounded if and only if*

$$g \in \bigcap_{i=0}^{m-1} V_{m-i}^{m-n-i-\frac{1}{p}+1}.$$

b) the operator $I^m_{g,\overline{a}} : H^p \to V_n$ *is compact if and only if*

$$g \in \bigcap_{i=0}^{m-1} V_{m-i,0}^{m-n-i-\frac{1}{p}+1}.$$

Remark 3.5. By choosing suitable parameters m, p, n and $\overrightarrow{g}$, the results obtained in this paper, can be stated for some well-known operators and spaces.

References

[1] E. Abbasi, The product-type operators from the Besov spaces into nth weighted type spaces. *Journal of Mathematical Extension,* 16(3): 1–14, 2022.

[2] E. Abbasi, S. Li and H. Vaezi, Weighted composition operators from the Bloch space to nth weighted-type spaces. *Turk. J. Math.,* 44(1): 108–117, 2020.

[3] E. Abbasi, Y. Liu and M. Hassanlou, Generalized Stevi-Sharma type operators from Hardy spaces into nth weighted type spaces. *Turk. J. Math.,* 45(4): 1543–1554, 2021.

[4] E. Abbasi, H. Vaezi and S. Li, Essential norm of weighted composition operators from H^∞ to nth weighted type spaces. *Mediterr. J. Math.,* 16: 133, 2019.

[5] H. Arroussi, H. Liu, C. Tong and Z. Yang, A new class of Carleson measures and integral operators on Bergman spaces. *Bull. Sci. Math.,* 197: 103531, 2024.

[6] N. Chalmoukis, Generalized integration operators on Hardy spaces. *Proc. Amer. Math. Soc.,* 148(8): 3325–3337, 2020.

[7] F. Colonna and N. Hmidouch, Weighted composition operators on iterated weighted-type Banach spaces of analytic functions. *Complex Anal. Oper. Theory,* 13: 1989–2016, 2019.

[8] J. Du, S. Li and D. Qu, The generalized Volterra integral operator and Toeplitz operator on weighted Bergman spaces. *Mediterr. J. Math.,* 19, 263, 2022.

[9] P. L. Duren, *Theory of H^p Spaces.* Academic Press, New York and London, 1970.

[10] C. Shen and S. Li, Volterra integral operators from Dirichlet type spaces into Hardy spaces. *Ric. mat.,* 72: 107–118, 2023.

[11] S. Stević, Composition operators from the Hardy space to the nth weighted-type space on the unit disk and the half-plane. *Appl. Math. Comput.,* 215(11): 3950–3955, 2010.

[12] S. Stevi, Weighted differentiation composition operators from H^∞ and Bloch spaces to nth weighted type spaces on the unit disk. *Appl. Math. Comput.,* 216: 3634–3641, 2010.

[13] R. Yang and S. Li, On a conjecture about generalized integration operators on Hardy spaces. arXiv preprint arXiv:2405.19372.

[14] K. Zhu, Bloch type space of analytic functions. *Rocky Mountain J. Math.,* 23: 1143–1177, 1993.

Ebrahim Abbasi
Department of Mathematics
Associate Professor of Mathematics
Mah. C., Islamic Azad University
Mahabad, Iran
E-mail: ebrahimabbasi81@aut.ac.ir, ebrahimabbasi81@gmail.com

Daryoush Molaei
Department of Mathematics
Assistant Professor of Mathematics
Ur. C., Islamic Azad University
Urmia, Iran
E-mail: daryoush.molaei@iau.ac.ir

Ali Ebrahimi
Department of Mathematics
Assistant Professor of Mathematics
Mah. C., Islamic Azad University
Mahabad, Iran
E-mail: ebrahimi.ali2719@aut.ac.ir, ebrahimi.ali2719@gmail.com