

## Tensor Product of Operator-Valued Frames in Hilbert C\*-Modules

**K. Musazadeh**

Islamic Azad University, Mahabad-Branch

**A. Khosravi**

Tarbiat Moallem University

**Abstract.** We show that the tensor product of two operator-valued frames for two Hilbert C\*-modules is an operator-valued frame for the tensor product of these Hilbert C\*-modules.

**AMS Subject Classification:** 94A12; 42C15; 68M10; 46C05.

**Keywords and Phrases:** Frames, operator-valued frames, tensor product, Hilbert C\*-modules.

### 1. Introduction

Frames on Hilbert C\*-modules have been defined for unital C\*-algebras by Frank and Larson [1] and investigated by many authors, see [2, 6, 11]. Recently, some generalizations of frames are proposed, for example, fusion frames, g-frames ([7]), operator-valued frames on Hilbert C\*-modules for a unital C\*-algebra ([4]). Furthermore, frames and bases in tensor products of Hilbert C\*-modules have been studied in [6]. For more details about the tensor product of Hilbert spaces and C\*-algebras we refer to [9]. We note that Hilbert C\*-modules are used in the study of locally compact quantum groups, completely positive maps between C\*-algebras, noncommutative geometry and K-theory. Also tensor product is useful in the approximation of multi-variate functions of combinations of univariate ones. In this section we recall some of the essential definitions and results which are needed in the sequel. For more details see [4].

Let  $\mathcal{A}$  be a  $C^*$ -algebra. We denote the Hilbert (right)  $C^*$ - $\mathcal{A}$ -module by  $\mathcal{H}_{\mathcal{A}}$ . The classic example of Hilbert (right)  $\mathcal{A}$ -module and the only one we will consider in this paper is the standard module  $\mathcal{H}_{\mathcal{A}} := \ell^2(\mathcal{A})$ , the space of all sequences  $\{a_i\}_{i \in I} \subset \mathcal{A}$  such that  $\sum_{i \in I} a_i^* a_i$  converges in norm to a positive element of  $\mathcal{A}$ .  $\ell^2(\mathcal{A})$  is endowed with the natural linear structure and right  $\mathcal{A}$ -multiplication, and with the  $\mathcal{A}$ -valued inner product defined by  $\langle \{a_i\}, \{b_i\} \rangle = \sum_{i \in I} a_i^* b_i$ , where the sum converges in norm by the Schwartz Inequality ([5]).

A map  $T$  from  $\mathcal{H}_{\mathcal{A}}$  to  $\mathcal{H}_{\mathcal{A}}$  is adjointable if there is a map  $T^* : \mathcal{H}_{\mathcal{A}} \rightarrow \mathcal{H}_{\mathcal{A}}$  such that  $\langle T^* \xi, \eta \rangle = \langle \xi, T \eta \rangle$  for all  $\xi, \eta \in \mathcal{H}_{\mathcal{A}}$ , ([4]). The collection of adjointable operators is denoted by  $\mathcal{B}(\mathcal{H}_{\mathcal{A}})$ . Then  $\mathcal{B}(\mathcal{H}_{\mathcal{A}})$  is a  $C^*$ -algebra ([5]). For each pair of elements  $\xi, \eta \in \mathcal{H}_{\mathcal{A}}$ , a bounded rank-one operator is defined by  $\theta_{\xi, \eta}(\zeta) = \xi \langle \eta, \zeta \rangle$ , for all  $\zeta \in \mathcal{H}_{\mathcal{A}}$ . The closed submodule of  $\mathcal{B}(\mathcal{H}_{\mathcal{A}})$  generated by rank-one operators is denoted by  $\mathcal{K}(\mathcal{H}_{\mathcal{A}})$ . When  $\mathcal{A} = \mathbb{C}$ , then  $\mathcal{H}_{\mathcal{A}} = \ell^2$ ,  $\mathcal{B}(\mathcal{H}_{\mathcal{A}}) = \mathcal{B}(\ell^2)$ , and  $\mathcal{K}(\mathcal{H}_{\mathcal{A}})$  coincides with the ideal  $\mathcal{K}$  of all compact operators on  $\ell^2$ .  $\mathcal{K}(\mathcal{H}_{\mathcal{A}})$  is always a closed ideal of  $\mathcal{B}(\mathcal{H}_{\mathcal{A}})$ . The analog of the strong operator topology on  $\mathcal{B}(\ell^2)$  is the *strict topology* on  $\mathcal{B}(\mathcal{H}_{\mathcal{A}})$  defined by

$$\mathcal{B}(\mathcal{H}_{\mathcal{A}}) \ni T_\lambda \rightarrow T \text{ strictly if } \|(T_\lambda - T)S\| \rightarrow 0 \text{ and } \|S(T_\lambda - T)\| \rightarrow 0, \forall S \in \mathcal{K}(\mathcal{H}_{\mathcal{A}}).$$

There is an alternate view of the objects  $\mathcal{B}(\mathcal{H}_{\mathcal{A}})$  and  $\mathcal{K}(\mathcal{H}_{\mathcal{A}})$ . Embed the tensor product  $\mathcal{A} \otimes \mathcal{K}$  into its Banach space second dual  $(\mathcal{A} \otimes \mathcal{K})^{**}$ , which, as is well known, is a  $W^*$ -algebra ([10]). The multiplier algebra of  $\mathcal{A} \otimes \mathcal{K}$ , is defined as the collection

$$\{T \in (\mathcal{A} \otimes \mathcal{K})^{**} : TS, ST \in \mathcal{A} \otimes \mathcal{K} \quad \forall S \in \mathcal{A} \otimes \mathcal{K}\}.$$

Equipped with the norm of  $(\mathcal{A} \otimes \mathcal{K})^{**}$ ,  $M(\mathcal{A} \otimes \mathcal{K})$  is a  $C^*$ -algebra. Assuming that  $\mathcal{A}$  is unital we apply the following two  $*$ -isomorphisms without further references:

$$\mathcal{B}(\mathcal{H}_{\mathcal{A}}) \cong M(\mathcal{A} \otimes \mathcal{K}) \quad \text{and} \quad \mathcal{K}(\mathcal{H}_{\mathcal{A}}) \cong \mathcal{A} \otimes \mathcal{K}.$$

The algebra  $\mathcal{B}(\mathcal{H}_{\mathcal{A}})$  is technically hard to work with, while  $M(\mathcal{A} \otimes \mathcal{K})$  is more accessible due to many established results. More details on the subject can be found in ([6, 8]). We denote the tensor product of two Hilbert  $C^*$ -modules  $\mathcal{H}_{\mathcal{A}}$  and  $\mathcal{H}_{\mathcal{B}}$  by  $\mathcal{H}_{\mathcal{A}} \otimes \mathcal{H}_{\mathcal{B}}$  which is a Hilbert  $\mathcal{A} \otimes \mathcal{B}$ -module, where  $\mathcal{A}$  and  $\mathcal{B}$  are  $C^*$ -algebras. See ([6, 8]).

## 2. Operator-Valued Frames on Hilbert C\*-Modules

We generalize an important result about the tensor product of frames on Hilbert C\*-modules to operator-valued frames. First, we recall some definitions.

**Definition 2.1.** *According to ([1]), a (vector) frame on the Hilbert C\*-module  $\mathcal{H}_A$  of a unital C\*-algebra  $\mathcal{A}$  is a collection of elements  $\{\xi_i\}_{i \in I}$  in  $\mathcal{H}_A$  for which there are two positive scalars  $a$  and  $b$  such that for all  $\xi \in \mathcal{H}_A$ ,*

$$a \langle \xi, \xi \rangle \leq \sum_{i \in I} \langle \xi, \xi_i \rangle \langle \xi_i, \xi \rangle \leq b \langle \xi, \xi \rangle,$$

where the convergence is in the norm of the C\*-algebra  $\mathcal{A}$ .

Let  $\eta \in \mathcal{H}_A$  be an arbitrary unital vector, i.e.,  $\langle \eta, \eta \rangle = Id$ , then by a result in ([4]),  $E_0 := \theta_{\eta, \eta} \in \mathcal{A} \otimes \mathcal{K}$  is a projection. Then  $\mathcal{H}_0 := E_0 \mathcal{H}_A$  is a submodule of  $\mathcal{H}_A$  and we can identify  $E_0 M(\mathcal{A} \otimes \mathcal{K})$  with  $B(\mathcal{H}_A, \mathcal{H}_0)$ , the set of linear bounded adjointable operators from  $\mathcal{H}_A$  to the submodule  $\mathcal{H}_0$ .

**Definition 2.2.** *Let  $\mathcal{A}$  be a unital C\*-algebra and  $I$  be a countable index set. Let  $E_0$  be a projection in  $M(\mathcal{A} \otimes \mathcal{K})$ . Denote by  $\mathcal{H}_0$  the submodule  $E_0 \mathcal{H}_A$  and identify  $B(\mathcal{H}_A, \mathcal{H}_0)$  with  $E_0 M(\mathcal{A} \otimes \mathcal{K})$ . A collection  $\{\Lambda_i \in B(\mathcal{H}_A, \mathcal{H}_0) : i \in I\}$  is called an operator-valued frame on  $\mathcal{H}_A$  with range in  $\mathcal{H}_0$  if the sum  $\sum_{i \in I} \Lambda_i^* \Lambda_i$  converges in the strict topology to a bounded invertible operator on  $\mathcal{H}_A$  denoted by  $D_\Lambda$ .  $\{\Lambda_i\}_{i \in I}$  is called a tight operator-valued (resp., Parseval operator-valued frame) if  $D_\Lambda = \lambda Id_{\mathcal{H}_A}$  for a positive number  $\lambda$  (resp.,  $D_\Lambda = Id_{\mathcal{H}_A}$ ), ([4]).*

Notice that if  $\{\Lambda_i\}_{i \in I}$  is an operator-valued frame with range in  $\mathcal{H}_0$ , then it is also an operator-valued frame with range in any larger submodule. For more details see ([4]). Most properties of frames on Hilbert spaces hold also for Hilbert C\*-modules. Also Kaftal, Larson and Zhang ([4]) showed that most results for operator-valued frames on Hilbert spaces are also true for operator-valued frames on Hilbert C\*-modules. In this section we have another result about this subject which

is a generalization of a result in ([6]).

**Lemma 2.3.** *Let  $\{T_\alpha\}_\alpha \subset B(\mathcal{H}_A)$  converges strictly to  $T$  and  $\{S_\beta\}_\beta \subset B(\mathcal{H}_A)$  converges strictly to  $S$ . Then  $\{T_\alpha \otimes S_\beta\}_{\alpha,\beta} \subset B(\mathcal{H}_A \otimes \mathcal{H}_A)$  converges strictly to  $T \otimes S$ .*

**Proof.** We first show that  $\mathcal{K}(\mathcal{H}_A) \otimes \mathcal{K}(\mathcal{H}_A)$  and  $\mathcal{K}(\mathcal{H}_A \otimes \mathcal{H}_A)$  are at least algebraically equivalent. For this, let  $\xi_1, \xi_2, \eta_1, \eta_2 \in \mathcal{H}_A$ . Then for all  $\zeta_1, \zeta_2 \in \mathcal{H}_A$  we have

$$\begin{aligned} (\theta_{\xi_1, \eta_1} \otimes \theta_{\xi_2, \eta_2})(\zeta_1 \otimes \zeta_2) &= \theta_{\xi_1, \eta_1}(\zeta_1) \otimes \theta_{\xi_2, \eta_2}(\zeta_2) = \xi_1 \langle \eta_1, \zeta_1 \rangle \otimes \xi_2 \langle \eta_2, \zeta_2 \rangle \\ &= (\xi_1 \otimes \xi_2)(\langle \eta_1, \zeta_1 \rangle \otimes \langle \eta_2, \zeta_2 \rangle) = (\xi_1 \otimes \xi_2) \langle \eta_1 \otimes \eta_2, \zeta_1 \otimes \zeta_2 \rangle \\ &= \theta_{\xi_1 \otimes \xi_2, \eta_1 \otimes \eta_2}(\zeta_1 \otimes \zeta_2). \end{aligned}$$

Therefore,  $\theta_{\xi_1, \eta_1} \otimes \theta_{\xi_2, \eta_2} = \theta_{\xi_1 \otimes \xi_2, \eta_1 \otimes \eta_2}$ . Since  $\mathcal{K}(\mathcal{H}_A) \otimes \mathcal{K}(\mathcal{H}_A)$  and  $\mathcal{K}(\mathcal{H}_A \otimes \mathcal{H}_A)$  are closed linear spans of these rank-one operators, the result follows. Now let  $U, V \in \mathcal{K}(\mathcal{H}_A)$ . Then  $U \otimes V \in \mathcal{K}(\mathcal{H}_A) \otimes \mathcal{K}(\mathcal{H}_A) = \mathcal{K}(\mathcal{H}_A \otimes \mathcal{H}_A)$ . So we have

$$\begin{aligned} &\|(T_\alpha \otimes S_\beta - T \otimes S)(U \otimes V)\| \\ &= \|(T_\alpha \otimes S_\beta - T_\alpha \otimes S + T_\alpha \otimes S - T \otimes S)(U \otimes V)\| \\ &= \|[T_\alpha \otimes (S_\beta - S) + (T_\alpha - T) \otimes S](U \otimes V)\| \\ &= \|[T_\alpha \otimes (S_\beta - S)](U \otimes V) + [(T_\alpha - T) \otimes S](U \otimes V)\| \\ &\leq \|T_\alpha U \otimes (S_\beta - S)V\| + \|(T_\alpha - T)U \otimes SV\| \\ &\leq \|T_\alpha U\| \|(S_\beta - S)V\| + \|(T_\alpha - T)U\| \|SV\|. \end{aligned}$$

Since  $\|(S_\beta - S)V\| \rightarrow 0$  and  $\|(T_\alpha - T)U\| \rightarrow 0$  by the hypothesis,

$$\|(T_\alpha \otimes S_\beta - T \otimes S)(U \otimes V)\| \rightarrow 0.$$

Similarly, we can show that  $\|(U \otimes V)(T_\alpha \otimes S_\beta - T \otimes S)\| \rightarrow 0$ , Hence the result follows.  $\square$

**Proposition 2.4.** *Let  $\{\Lambda_i\}_{i \in I}$  and  $\{\Gamma_i\}_{i \in I}$  be operator-valued frames in  $B(\mathcal{H}_A, \mathcal{H}_0)$ . Then  $\{\Lambda_i \otimes \Gamma_j\}_{i,j}$  is an operator-valued frame in  $B(\mathcal{H}_A \otimes \mathcal{H}_A, \mathcal{H}_0 \otimes \mathcal{H}_0)$ .*

**Proof.** Suppose that  $\{\Lambda_i\}_{i \in I}$  and  $\{\Gamma_i\}_{i \in I}$  are operator-valued frames in  $B(\mathcal{H}_A, \mathcal{H}_0)$ . Then we have

$$\begin{aligned} \sum_{i,j} (\Lambda_i \otimes \Gamma_j)^* (\Lambda_i \otimes \Gamma_j) &= \sum_{i,j} (\Lambda_i^* \otimes \Gamma_j^*) (\Lambda_i \otimes \Gamma_j) = \sum_{i,j} (\Lambda_i^* \Lambda_i \otimes \Gamma_j^* \Gamma_j) \\ &= \sum_{i \in I} \Lambda_i^* \Lambda_i \otimes \sum_{j \in I} \Gamma_j^* \Gamma_j. \end{aligned}$$

But  $\sum_{i \in I} \Lambda_i^* \Lambda_i$  converges strictly to  $S_\Lambda$  and  $\sum_{j \in I} \Gamma_j^* \Gamma_j$  converges strictly to  $S_\Gamma$  by the definition. Hence by the above lemma  $\sum_{i \in I} \Lambda_i^* \Lambda_i \otimes \sum_{j \in I} \Gamma_j^* \Gamma_j$  converges strictly to  $S_\Lambda \otimes S_\Gamma$ , which is a bounded invertible operator on  $\mathcal{H}_A \otimes \mathcal{H}_A$ .  $\square$

Since by Theorem 1.4. of ([4]) a collection  $\{\xi_i\}_{i \in I}$  in  $\mathcal{H}_A$  is a frame if and only if  $\{A_i \in B(\mathcal{H}_A, \mathcal{H}_0) : i \in I\}$  is an operator-valued frame in  $\mathcal{H}_A$ , where  $A_i(\xi) = \langle \xi, \xi_i \rangle$ , as a particular case of Proposition 2.4. we get a result proved in ([6]).

**Corollary 2.5.** *Let  $\{f_i\}_{i \in I}$  and  $\{g_i\}_{i \in I}$  be standard frames for  $\mathcal{H}_A$  and  $\mathcal{H}_B$ , respectively. Then  $\{f_i \otimes g_j\}_{i,j \in I}$  is a standard frame for  $\mathcal{H}_A \otimes \mathcal{H}_B$ .*

**Proof.** See ([6]).  $\square$

Generalized frames in Hilbert C\*-modules is another aspect of operator-valued frames in Hilbert C\*-modules which is introduced in ([7]).

### Acknowledgement

The authors express their gratitude to the referee for his careful reading and for several valuable points which improved the manuscript.

## References

- [1] M. Frank and D. Larson, Frames in Hilbert  $C^*$ -modules and  $C^*$ -algebras, *J. Operator Theory*, 48 (2002), 273-314.
- [2] D. Han, W. Jing, and R. M. Mohapatra, Structured Parseval frames in Hilbert  $C^*$ -modules, (*preprint*).
- [3] V. Kaftal, D. Larson, and S. Zhang, Operator valued frames on  $C^*$ -modules, *Contemporary Mathematics*, 2007.
- [4] V. Kaftal, D. Larson, and S. Zhang, Operator valued frames, *Trans. Amer. Math. Soc.* 361 (2009), 6349-6385.
- [5] G. G. Kasparov, Hilbert  $C^*$ -modules: theorems of Stinespring and Voiculescu, *J. Operator Theory*, 3 (1980), 133-150.
- [6] A. Khosravi and B. Khosravi, Frames and bases in tensor products of Hilbert spaces and Hilbert  $C^*$ -modules, *Proc. Indian Acad. Sci. (Math. Sci.)*, 117 (1) (2007), 1-12.
- [7] A. Khosravi and B. Khosravi, Fusion frames and g-frames in Hilbert  $C^*$ -modules, *International Journal of Wavelets, Multiresolution and Information Processing*, 6 (3) (2008), 433-446.
- [8] E. C. Lance, *Hilbert  $C^*$ -modules-a toolkit for operator algebraists*, London Math. Soc. Lecture Note Ser., (Cambridge, England: Cambridge Univ. Press) (1995), vol. 210.
- [9] G. J. Murphy,  *$C^*$ -algebras and operator theory*, Academic Press, San Diego, California, 1990.
- [10] G. K. Pedersen,  *$C^*$ -algebras and their automorphism groups*, Academic Press, London, New York, San Francisco, 1979.
- [11] P. Wood, Wavelets and projective Hilbert modules , (*preprint*).

**Kamran Musazadeh**

Department of Mathematics  
Islamic Azad University, Mahabad Branch  
Mahabad, Iran.  
E-mail: kamran\_ms2004@yahoo.com

**Amir Khosravi**

Faculty of Mathematical Sciences and Computer  
Tarbiat Moallem University  
Tehran, Iran.  
E-mail: khosravi\_amir@yahoo.com