

A New Three-step Iterative Method for Solving Nonlinear Equations

M. Matin Far*

University of Mazandaran

M. Aminzadeh

University of Mazandaran

S. Asadpour

University of Mazandaran

Abstract. In this paper, a new three-step iterative method for finding a simple root of the nonlinear equation of $f(x) = 0$ will be introduced. This method is based on the two-step method of [C. Chun, Y. Ham, Some fourth-order modifications of Newton's method, Appl. Math. Comput. 197 (2008) 654-658]. The new method requires three evaluations of the function and two of its first-derivative. We will prove that the order of convergence of the new method and its efficiency index will respectively be 8, and 1.5157. Some numerical experiments are given to illustrate the performance of the three-step iterative method.

AMS Subject Classification: 65B99; 65N99.

Keywords and Phrases: Nonlinear equation, iterative method, three-step iterative method, convergence order, efficiency index.

1. Introduction

In recent years, attempts have been made to increase the order of convergence, and improve the efficiency index in iterative methods, examples of which can be found in [3–10]. Most of the mentioned papers have used

Received: December 2011; Final Revision: January 2012

*Corresponding author

the Newton's method to calculate the final approximation of the exact solution of the equation $f(x) = 0$. The calculation of the derivative in the third step is the fundamental drawback of these methods, which in turn decreases their efficiency index. However, the efficiency index of these methods can be increased by using approximation for calculating the derivatives. In this paper, a new three-step iterative method for solving nonlinear equations will be introduced. This method, using Stefensen's method has solved the problem of the calculation of derivative in the third step. This method also has an order of convergence of 8, and an efficiency index of 1.5157.

2. Development of Method and Convergence Analysis

In this section, we will first describe the ideas of the iterative method proposed by C. Chun and Y. Ham in [2], and then will give a new three-step iterative method and its convergence analysis.

To solve nonlinear equations, iterative methods such as Newton's method are usually used. Throughout this paper we will consider iterative methods to find a simple root, i.e., $f(\alpha) = 0$ and $f'(\alpha) \neq 0$, of a nonlinear equation $f(x) = 0$ that uses f and f' but not the higher derivatives of f . Newton's method for the calculation of a simple root is probably the most widely used iterative methods defined by

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}. \quad (1)$$

It is well known ([12]) that this method is quadratically convergent. We will here consider Newton's method as the first step, and hence have:

$$y_n = x_n - \frac{f(x_n)}{f'(x_n)}. \quad (2)$$

Now, we will consider the following two-step methods: double-Newton method with fourth-order convergence ([12]) given by

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} - \frac{f(y_n)}{f'(y_n)}, \quad (3)$$

the fourth-order method [1] given by

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} - \left[1 + 2 \frac{f(y_n)}{f(x_n)} + \frac{f(y_n)^2}{f(x_n)^2} \right] \frac{f(y_n)}{f'(x_n)}, \quad (4)$$

and King's fourth-order family of methods [7] given by

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} - \frac{f(x_n) + \beta f(y_n)}{f(x_n) + (\beta - 2)f(y_n)} \frac{f(y_n)}{f'(x_n)}, \quad (5)$$

where $\beta \in \mathbb{R}$.

Approximately equaling the correction terms of the methods (3) and (4), we can obtain an approximation to $f'(y_n)$

$$f'(y_n) \approx \frac{f'(x_n)f(x_n)^2}{f(x_n)^2 + 2f(x_n)f(y_n) + f(y_n)^2}. \quad (6)$$

Approximately equaling the correction terms of the methods (3) and (5), we can obtain an approximation to $f'(y_n)$

$$f'(y_n) \approx \frac{f'(x_n)[f(x_n) + (\beta - 2)f(y_n)]}{f(x_n) + \beta f(y_n)}. \quad (7)$$

Now, we apply the respective approximation (6) and (7) to any other iterative method whose order is at least four and which also depend on $f'(y_n)$, for example, the fifth-order method proposed in [5] given by

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} - \frac{f'(y_n) + 3f'(x_n)}{5f'(y_n) - f'(x_n)} \frac{f(y_n)}{f'(x_n)}. \quad (8)$$

Per iteration this method requires two evaluations of the function and two evaluations of its first-derivative, so its efficiency index equals to $5^{1/4} \approx 1.495$ if we consider the definition of efficiency index [7] as $p^{1/m}$, where p is the order of the method and m is the number of functional evaluations per iteration required by the method. Using (7) in (8), we obtain the family of methods

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} - \frac{2f(x_n) + (2\beta - 1)f(y_n)}{2f(x_n) + (2\beta - 5)f(y_n)} \frac{f(y_n)}{f'(x_n)}, \quad (9)$$

where $\beta \in \mathfrak{R}$. The family (9) includes, as particular cases, the following ones: For $\beta = 0$, we obtain a fourth-order method which is the second step of our three-step method:

$$z_n = x_n - \frac{f(x_n)}{f'(x_n)} - \frac{2f(x_n) - f(y_n)}{2f(x_n) - 5f(y_n)} \frac{f(y_n)}{f'(x_n)}, \quad (10)$$

Now we will introduce Steffensen's method, by putting z_n in which, our new three-step method will be created. This method has the second order of convergence, and contrary to Newton's method, does not require the calculation of the function's derivative. In this method, $f'(x_n)$, in the procedure of Newton's method will be substituted by the function

$$g(x_n) = \frac{f(x_n + f(x_n)) - f(x_n)}{f(x_n)}, \quad (11)$$

and one can obtain the iterative procedure of Steffensen as follows:

$$x_{n+1} = x_n - \frac{f(x_n)}{g(x_n)} = x_n - \frac{(f(x_n))^2}{f(x_n + f(x_n)) - f(x_n)} \quad n = 0, 1, 2, \dots \quad (12)$$

Now, we will introduce our new three-step method:

$$\begin{aligned} y_n &= x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n &= x_n - \frac{f(x_n)}{f'(x_n)} - \frac{2f(x_n) - f(y_n)}{2f(x_n) - 5f(y_n)} \frac{f(y_n)}{f'(x_n)}, \\ x_{n+1} &= z_n - \frac{(f(z_n))^2}{f(z_n + f(z_n)) - f(z_n)} \quad n = 0, 1, 2, \dots \end{aligned} \quad (13)$$

For the methods defined by (13), we have

Theorem : *Let $\alpha \in I$ be a simple zero of a sufficiently differentiable function $f : I \rightarrow \mathbb{R}$ for an open interval I . If x_0 is sufficiently close to α , then the three-step iterative method (13) has eighth-order convergence and satisfies the following error equation :*

$$e_{n+1} = -2(c_2^2 c_3^2) e_n^8 + O(e_n^9), \quad (14)$$

where $e_n = x_n - \alpha$ for $n = 0, 1, 2, \dots$, and $c_n = \frac{f^{(n)}(\alpha)}{n! f'(\alpha)}$.

Proof. Since f is sufficiently differentiable, by expanding $f(x_n)$ and $f'(x_n)$ about α , one obtains

$$\begin{aligned} f(x_n) &= f'(\alpha)(e_n + c_2 e_n^2 + c_3 e_n^3 + c_4 e_n^4 + c_5 e_n^5 \\ &\quad + c_6 e_n^6 + c_7 e_n^7 + c_8 e_n^8 + O(e_n^9)), \\ f'(x_n) &= f'(\alpha)(1 + 2c_2 e_n + 3c_3 e_n^2 + 4c_4 e_n^3 + 5c_5 e_n^4 + 6c_6 e_n^5 \\ &\quad + 7c_7 e_n^6 + 8c_8 e_n^7 + O(e_n^8)). \end{aligned} \quad (15)$$

where $c_k = \frac{1}{k!} \frac{f^{(k)}(\alpha)}{f'(\alpha)}$ for $k = 2, 3, \dots$. From (13), and (15), using the

Maple software one obtains

$$\begin{aligned} y_n &= \alpha + c_2 e_n^2 - 2(c_2^2 - c_3)e_n^3 - (7c_2 c_3 - 3c_4 - 4c_2^3)e_n^4 \\ &\quad - (8c_2^4 - 20c_3 c_2^2 + 6c_3^2 + 10c_4 c_2 - 4c_5)e_n^5 + (-17c_4 c_3 \\ &\quad + 28c_4 c_2^2 - 13c_2 c_5 + 5c_6 + 33c_2 c_3^2 - 52c_3 c_2^3 + 16c_2^5)e_n^6 \\ &\quad + (92c_3 c_2 c_4 - 22c_3 c_5 + 18c_3^3 - 126c_3^2 c_2^2 + 128c_3 c_4^2 \\ &\quad - 12c_4^2 - 72c_4 c_2^3 + 36c_5 c_2^2 + 6c_7 - 16c_2 c_6 - 32c_2^6)e_n^7 \\ &\quad + (7c_8 + 118c_5 c_2 c_3 - 348c_4 c_3 c_2^2 - 19c_2 c_7 + 64c_2^7 \\ &\quad + 64c_2 c_4^2 - 31c_4 c_5 + 75c_4 c_3^2 + 176c_4 c_2^4 - 92c_5 c_2^3 - 27c_6 c_3 \\ &\quad + 44c_6 c_2^2 - 135c_2 c_3^3 + 408c_3^2 c_2^3 - 304c_3 c_2^5)e_n^8 + O(e_n^9). \end{aligned} \quad (16)$$

Expanding $f(y_n)$ about α and using (16),

$$\begin{aligned} f(y_n) &= f'(\alpha)(c_2 e_n^2 - 2(c_2^2 - c_3)e_n^3 - (7c_2 c_3 - 3c_4 - 5c_2^3)e_n^4 \\ &\quad + (4c_5 - 12c_2^4 - 6c_3^2 - 10c_2 c_4 + 24c_3 c_2^2)e_n^5 + (-13c_2 c_5 \\ &\quad - 73c_3 c_2^3 + 34c_4 c_2^2 + 5c_6 - 17c_4 c_3 + 37c_2 c_3^2 + 28c_2^5)e_n^6 \\ &\quad + (44c_5 c_2^2 + 104c_3 c_2 c_4 - 22c_3 c_5 + 18c_3^3 + 6c_7 + 206c_3 c_2^4 \\ &\quad - 104c_4 c_2^3 - 160c_3^2 c_2^2 - 64c_2^6 - 12c_4^2 - 16c_2 c_6)e_n^7 + (75c_4 c_3^2 \\ &\quad + 134c_5 c_2 c_3 + 73c_2 c_4^2 + 34c_6 c_2^2 - 552c_3 c_2^5 - 147c_2 c_3^3 + 7c_8 \\ &\quad - 19c_2 c_7 + 297c_4 c_2^4 - 134c_5 c_2^3 + 582c_3^2 c_2^3 + 144c_2^7 - 27c_6 c_3 \\ &\quad - 455c_4 c_3 c_2^2 - 31c_4 c_5)e_n^8 + O(e_n^9)). \end{aligned} \quad (17)$$

From (13), one obtains

$$\begin{aligned} z_n &= \alpha - c_2 c_3 e_n^4 + ((3/2)c_2^4 - 2c_3^2 + 2c_3 c_2^2 - 2c_2 c_4)e_n^5 \\ &\quad + (8c_3 c_2^3 + 6c_2 c_3^2 - 7c_4 c_3 - 3c_2 c_5 - (35/4)c_2^5 + 3c_4 c_2^2)e_n^6 \\ &\quad + (4c_5 c_2^2 + (255/8)c_2^6 + 14c_4 c_2^3 + 16c_3 c_2 c_4 - 10c_3 c_5 \\ &\quad + 4c_3^3 + 16c_3^2 c_2^2 - 4c_2 c_6 - (135/2)c_3 c_2^4 - 6c_4^2)e_n^7 + (19c_5 c_2^3 \\ &\quad - 5c_2 c_7 + 14c_4 c_3^2 + 20c_5 c_2 c_3 - 13c_6 c_3 - (413/4)c_4 c_2^4 + 61c_4 c_3 c_2^2 \\ &\quad + 13c_2 c_3^3 - (1515/16)c_2^7 + 10c_2 c_4^2 - 199c_3^2 c_2^3 - 17c_4 c_5 \\ &\quad + (573/2)c_3 c_2^5 + 25c_6 c_2^2)e_n^8 + O(e_n^9). \end{aligned} \quad (18)$$

Expanding $f(z_n)$ about α and using (18),

$$\begin{aligned}
f(z_n) = & f'(\alpha)(-c_2c_3e_n^4 + ((3/2)c_2^4 + 2c_3c_2^2 - 2c_3^2 - 2c_2c_4)e_n^5 \\
& + (-35/4)c_2^5 - 7c_4c_3 - 3c_2c_5 + 8c_3c_2^3 + 6c_2c_3^2 + 3c_4c_2^2)e_n^6 \\
& + (4c - 3^3 + (255/8)c_2^6 - 6c_4^2 + 16c_3c_2c_4 - (135/2)c_3c_2^4 \\
& - 10c_3c_5 + 14c_4c_2^3 + 16c_3^2c_2^2 - 4c_2c_6 + 4c_5c_2^2)e_n^7 + (14c_4c_3^2 \\
& + (573/2)c_3c_2^5 - 5c_2c_7 - 17c_4c_5 + 25c_6c_2^2 - 13c_6c_3 \\
& - (413/4)c_4c_2^4 + c_2^2c_3^2 - 199c_3^2c_2^3 - (1515/16)c_2^7 + 19c_5c_2^3 \\
& + 13c_2c_3^3 + 10c_2c_4^2 + 61c_4c_3c_2^2 + 20c_5c_2c_3)e_n^8 + O(e_n^9)). \tag{19}
\end{aligned}$$

Expanding $f(z_n + f(z_n))$ about α and using (18) and (19),

$$\begin{aligned}
f(z_n + f(z_n)) = & f'(\alpha)(-2c_2c_3e_n^4 + (3c_2^4 - 4c_2c_4 + 4c_3c_2^2 - 4c_3^2)e_n^5 \\
& + (-14c_4c_3 - 6c_2c_5 + 16c_3c_2^3 + 6c_4c_2^2 + 12c_2c_3^2 \\
& - (35/2)c_2^5)e_n^6 + (28c_4c_2^3 + 32c_3^2c_2^2 + 32c_3c_2c_4 - 135c_3c_2^4 \\
& - 20c_3c_5 + 8c_5c_2^2 - 8c_2c_6 + 8c_3^3 + (255/4)c_2^6 - 12c_4^2)e_n^7 \\
& + (26c_2c_3^3 - 26c_6c_3 + 50c_6c_2^2 + 40c_5c_2c_3 + 122c_4c_3c_2^2 \\
& + 20c_2c_4^2 + 573c_3c_2^5 + 28c_4c_3^2 - 398c_3^2c_2^3 + 38c_5c_2^3 \\
& - 34c_4c_5 - 10c_2c_7 - (1515/8)c_2^7 + c_2^2c_3^2 \\
& - (413/2)c_4c_2^4)e_n^8 + O(e_n^9)). \tag{20}
\end{aligned}$$

Using (19),

$$\begin{aligned}
(f(z_n))^2 = & (f'(\alpha))^2((c_2^2c_3^2)e_n^8 + (4c_2c_3^3 - 4c_3^3c_2^2 + 4c_4c_3c_2^2 \\
& - 3c_2^5c_3)e_n^9 + (6c_2^2c_5c_3 + (47/2)c_3c_2^6 - 6c_2^5c_4 \\
& - 14c_3c_2^3c_4 + (9/4)c_2^8 - 18c_3^2c_2^4 + 22c_4c_3^2c_2 - 20c_3^3c_2^2 \\
& + 4c_2^2c_4^2 + 4c_3^4)e_n^{10} + O(e_n^{11})). \tag{21}
\end{aligned}$$

From (13), (20) and (21), one obtains

$$x_{n+1} = \alpha - (2c_2^2c_3^2)e_n^8 + O(e_n^9), \tag{22}$$

which implies that $e_{n+1} = (-2c_2^2c_3^2)e_n^8 + O(e_n^9)$.

Since in each iteration our new method requires three functional evaluations, and two functional evaluation of its derivative; and since it has an order of convergence of 8, its efficiency index will equal : $8^{1/5} \approx 1.5157$.

3. Numerical Experiments

We present some numerical results to illustrate the efficiency of the three-step iterative method proposed in this paper.

We compare (13) (MAA means: M:Matinfar, A:Asadpour, A:Aminzadeh) with the existing three-step methods in [13] which are respectively represented by XIA 1, XIA 2 and XIA 3.

XIA 1 is :

$$\begin{aligned} y_n &= x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n &= x_n - \frac{f(x_n)}{f'(x_n)} - \frac{f(x_n) - f(y_n)}{f(x_n) - 2f(y_n)}, \\ x_{n+1} &= z_n - \frac{f(z_n)}{f'(z_n)} \left[1/2 + \frac{5f(x_n)^2 + 8f(x_n)f(y_n) + 2f(y_n)^2}{5f(x_n)^2 - 12f(x_n)f(y_n)} (1/2 + \frac{f(z_n)}{f(y_n)}) \right]. \end{aligned}$$

XIA 2 is :

$$\begin{aligned} y_n &= x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n &= x_n - \frac{f(x_n)}{f'(x_n)} - \frac{f(x_n) - f(y_n)}{f(x_n) - 2f(y_n)}, \\ x_{n+1} &= z_n - \frac{f(z_n)}{f'(z_n)} \left[\frac{5f(x_n)^2 - 2f(x_n)f(y_n) + f(y_n)^2}{5f(x_n)^2 - 12f(x_n)f(y_n)} + (1 + 4\frac{f(y_n)}{f(x_n)})(\frac{f(z_n)}{f(y_n)}) \right]. \end{aligned}$$

XIA 3 is :

$$\begin{aligned} y_n &= x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n &= x_n - \frac{f(x_n)}{f'(x_n)} - \frac{4f(x_n)^2 - 5f(x_n)f(y_n) - f(y_n)^2}{4f(x_n)^2 - 9f(x_n)f(y_n)}, \\ x_{n+1} &= z_n - \frac{f(z_n)}{f'(z_n)} \left[1 + 4\frac{f(z_n)}{f(x_n)} \right] \left[\frac{8f(y_n)}{4f(x_n) - 11f(y_n)} + 1 + \frac{f(z_n)}{f(y_n)} \right]. \end{aligned}$$

The iteration methods above will be stopped when the following stopping criteria is satisfied:

$$\left| f(x_k) \right| < \epsilon.$$

where ϵ represents tolerance. The test functions to be used are defined as follows:

$$\begin{aligned} f_1(x) &= x^3 + 4x^2 - 15, & x_* &= 1.63198055660636, \\ f_2(x) &= xe^{x^2} - \sin^2(x) + 3\cos(x) + 5, & x_* &= -1.207647827130919, \\ f_3(x) &= 10xe^{-x^2} - 1, & x_* &= 1.67963061042845, \\ f_4(x) &= \sin^2(x) - x^2 + 1, & x_* &= 1.4044916482153411, \\ f_5(x) &= x^5 + x^4 + 4x^2 - 15, & x_* &\approx 1.347. \end{aligned}$$

where x_* is the exact root. The absolute values of the function ($|f(x_n)|$) and ($|x_n - x_*|$), are shown in Table 1.

Table 1: Comparison of iterative methods.

Function	Method	$ x_k - x_* $	$f(x_k)$
$f_1(x), x_0 = 2$	MAA	2.489597032973023e-007	3.552713678800501e-015
	XIA 1	2.489597032973023e-007	3.552713678800501e-015
	XIA 2	2.489597035193469e-0074	3.552713678800501e-015
	XIA 3	2.489597035193469e-007	3.552713678800501e-015
$f_2(x), x_0 = -1$	MAA	0	2.664535259100376e-015
	XIA 1	6.661338147750939e-016	1.509903313490213e-014
	XIA 2	0	2.664535259100376e-015
	XIA 3	1.798561299892754e-014	3.632649736573512e-013
$f_3(x), x_0 = 1.5$	MAA	0	2.220446049250313e-016
	XIA 1	0	2.220446049250313e-016
	XIA 2	0	2.220446049250313e-016
	XIA 3	0	2.220446049250313e-016
$f_4(x), x_0 = 1.5$	MAA	0	3.330669073875470e-016
	XIA 1	0	3.330669073875470e-016
	XIA 2	2.220446049250313e-016	4.440892098500626e-016
	XIA 3	2.220446049250313e-016	4.440892098500626e-016
$f_5(x), x_0 = 1.2$	MAA	4.280989683049796e-004	1.776356839400251e-015
	XIA 1	4.280989683049796e-004	1.776356839400251e-015
	XIA 2	4.280989683049796e-004	1.776356839400251e-015
	XIA 3	4.280989683049796e-004	1.776356839400251e-015

All numerical tests have been written in Matlab. The numerical results imply that our three-step method has a good performance, and despite being simple and requiring a small number of calculations compared to other existing methods, delivers good numerical results.

4. Conclusions

In this paper, we proposed a new three-step iterative method for solving nonlinear equations by using Steffensen's method in the third step. We showed that the new three-step iterative method has eighth-order convergence. Numerical experiments also show that the numerical results of our new three-step method, in equal iterations, resemble the results of other existing three-step methods with eighth-order convergence. However, our new method relatively requires a smaller number of calculations and does not require the calculation of derivative in the third step.

References

- [1] C. Chun, Some fourth-order iterative methods for solving nonlinear equations, *Applied Mathematics and Computation*, doi:10.1016/j.amc.2007.04.105.
- [2] C. Chun, Y. Ham, Some fourth-order modifications of Newton's method, *Applied Mathematics and Computation*, 197 (2008), 654-658.
- [3] A. Cordero, J. L. Hueso, E. Martinez, and J. R. Torregrosa, A family of iterative methods with sixth and seventh order convergence for nonlinear equations, *Mathematical and Computer Modelling*, 52 (2010), 1490-1496.
- [4] A. Cordero, J. L. Hueso, E. Martinez, and J. R. Torregrosa, *A modified Newton Jarrat's composition*, *Numerical Algorithm*, doi:10.1007/s11075-009-9359-z.
- [5] Y. Ham, C. Chun, *A fifth-order iterative method for solving nonlinear equation*, Adoi:10.1016/ j.amc.2007.04.005.
- [6] Y. Ham, C. Chun, and S. G. Lee, Some higher-order modifications of Newton's method for solving nonlinear equations, *Journal of Computational and Applied Mathematics*, 222 (2008), 477-486.
- [7] R. King, A family of fourth-order methods for nonlinear equations, *SIAM J. Numer. Anal.*, 10 (5) (1973), 876-879.
- [8] J. Kou, Y. Li, and X. Wang, Some variants of Ostrowski's method with seventh-order convergence, *Journal of Computational and Applied Mathematics*, 209 (2007), 153-159.
- [9] L. Liu and X. Wang, Eighth-order methods with high efficiency index for solving nonlinear equations, *Applied Mathematics and Computation*, 215 (2010), 3449-3454.
- [10] N. A. Mir and T. Zaman, Some quadrature based three-step iterative methods for non-linear equations, *Applied Mathematics and Computation*, 193 (2007), 366-373.
- [11] S. K. Parhi and D. K. Gupta, A six order method for nonlinear equations, *Applied Mathematics and Computation*, 203 (2008), 50-55.
- [12] J. F. Traub, *Iterative Methods for the Solution of Equations*, Chelsea Publishing Company, New York, 1982.

- [13] X. Wang and L. Liu, New eighth-order iterative methods for solving non-linear equations, *Computational and Applied Mathematics*, 234 (2010), 1611-1620.
- [14] X. Wang and L. Liu, Two new families of sixth-order methods for solving non-linear equations, *Applied Mathematics and Computation*, 213 (2009), 73-78.
- [15] S. Weerakoon and T. G. I. Fernando, A variant of Newton's method with accelerated third-order convergence, *Applied Mathematics Letters*, 13 (2000), 87-93.

Mashallah Matin Far

Department of Mathematics and Computer Science

Assistant Professor of Mathematics

University of Mazandaran

P.O.Box 47415-95447

Babolsar, Iran

E-mail: m.matinfar@umz.ac.ir

Mohammad Aminzadeh

Department of Mathematics

M.Sc. Student

Science of Mathematics Faculty

University of Mazandaran

P.O.Box 47415-95447

Babolsar, Iran

E-mail: m.aminzadeh@stu.umz.ac.ir

Sasan Asadpour

Department of Mathematics

M.Sc. Student

Science of Mathematics Faculty

University of Mazandaran

P.O.Box 47415-95447

Babolsar, Iran

E-mail: sasan.a.g@gmail.com